



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A THEOREM CONCERNING EQUAL RATIOS

BY J. L. COOLIDGE

THE following simple theorem is not included in the common textbooks on the Theory of Numbers, but it is altogether likely that it has been published before. The writer would be grateful to any reader of THE ANNALS who should tell him where a demonstration has already appeared.

THEOREM: If a set of ratios between positive integers are equal to one another, then all are equal to the ratio of the greatest common divisor of the numerators to that of the denominators, and also to the ratio of the least common multiple of the numerators to that of the denominators.

To prove the theorem, let

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} \cdots = \frac{a_n}{b_n} = \frac{p}{q}$$

where p and q are any two positive integers in the same ratio as the positive integers, a_i and b_i where $i = 1, 2, \dots n$. We have then n equations

$$qa_i = pb_i.$$

The greatest common divisor of the first members of these equations must be equal to that of the second members, so that if a be the greatest common divisor of the a_i 's and b that of the b_i 's, then, since qa and pb are the greatest common divisors of the first and second members respectively, we have

$$qa = pb \quad \text{or} \quad \frac{a}{b} = \frac{p}{q}.$$

If, further, we put

$$a_i = r_i a \quad b_i = s_i b$$

then, since

$$\frac{a_i}{b_i} = \frac{a}{b}$$

we have $r_i = s_i$.

Now if A and B are the least common multiples of the a_i 's and b_i 's respectively, and if R and S are the least common multiples of the r_i 's and s_i 's respectively, we have

$$A = Ra \qquad B = Sb$$

and since, for all values of i , $r_i = s_i$, $R = S$, and consequently

$$\frac{A}{B} = \frac{a}{b} = \frac{p}{q}$$

CAMBRIDGE, MASS.,

JUNE, 1908.